

Retrieving Liquid Cloud Droplet Size Distribution using the Cloudbow Geometric Retrieval Algorithm

Rachel E. Smith^{a,b,c,*}, *, Xiaoguang Xu^{a,b,c}, Brent A. McBride^{a,b,c}, and J. Vanderlei Martins^{a,b,c}

^a Department of Physics, University of Maryland Baltimore County, Baltimore, Maryland, USA

^b Goddard Earth Sciences Technology and Research (GESTAR)II, Baltimore, Maryland, USA

^c Earth and Space Institute, University of Maryland Baltimore County, Baltimore, Maryland, USA

* Presenting author (rsmith12@umbc.edu)

Liquid cloud microphysics modulates the amount of short-wave radiation reflected back to space and the amount of long-wave radiation trapped close to Earth's surface [1]. A two-parameter gamma distribution (DSD) consisting of cloud droplet effective radius (CDR) and cloud droplet effective variance (CDV) is commonly used to represent liquid clouds. CDV is the width of the DSD and as such, it contains information on cloud evolution and radiative processes [2]. Previous works have shown that polarimetric measurements are sensitive to both DSD parameters simultaneously, and that CDR modulates the scattering angle of the primary and supernumerary cloudbow peaks while CDV varies the amplitude of the supernumerary bows [2]. The geometric parameters of the polarized cloudbow contain the first-order cloud DSD information which we have used to develop a Lookup Table (LUT). By correcting for atmospheric transmittance and geometrical dependence, we have developed a LUT of polarized reflectances which are applicable to a wide range of viewing geometries. This cloudbow geometric retrieval algorithm allows for the efficient retrieval of the liquid cloud DSD from multiangular polarimetric measurements such as those from the Hyper-Angular Rainbow Polarimeter (HARP) suite of instruments. The Hyper-Angular Rainbow Polarimeter 2 (HARP2) aboard NASA's Plankton, Aerosol, Cloud ocean Ecosystem (PACE) satellite, is a wide field-of-view instruments capable of measuring polarized radiance at ~ 2.5 km resolution near nadir and up to 60 viewing angles in the 670 nm spectral band [3]. In this work we will demonstrate global retrievals of liquid cloud droplet size distribution using the cloudbow geometric retrieval algorithm on HARP2 data and compare the retrievals with HARP2 operational cloud products.

References

- [1] Ramanathan, V., Cess, R. D., Harrison, E. F., Minnis, P., Barkstrom, B. R., Ahmad, E., & Hartmann, D.: Cloud-radiative forcing and climate: Results from the Earth Radiation Budget Experiment. *Science*, 243(4887), 57–63, <https://doi.org/10.1126/science.243.4887.57>, 1989.
- [2] Breon, F. M. and Goloub, P.: Cloud droplet effective radius from spaceborne polarization measurements, *Geophys. Res. Lett.*, 25, 1879–1882, <https://doi.org/10.1029/98gl01221>, 1998.
- [3] Martins J.V. et al., First Results and On-Orbit Performance of the Hyper-Angular Rainbow Polarimeter (HARP2) on the PACE Satellite, SPIE

Preferred mode of presentation: Oral/Poster